

CIVIL ENGINEERING

Engineering Mechanics



Comprehensive Theory
with Solved Examples and Practice Questions



MADE EASY Publications Pvt. Ltd.

Corporate Office: 44-A/4, Kalu Sarai (Near Hauz Khas Metro Station), New Delhi-110016 | **Ph. :** 9021300500

Email : infomep@madeeasy.in | **Web :** www.madeeasypublications.org

Engineering Mechanics

Copyright © by MADE EASY Publications Pvt. Ltd.
All rights are reserved. No part of this publication may be reproduced, stored in or introduced into a retrieval system, or transmitted in any form or by any means (electronic, mechanical, photo-copying, recording or otherwise), without the prior written permission of the above mentioned publisher of this book.



MADE EASY Publications Pvt. Ltd. has taken due care in collecting the data and providing the solutions, before publishing this book. In spite of this, if any inaccuracy or printing error occurs then **MADE EASY Publications Pvt. Ltd.** owes no responsibility. We will be grateful if you could point out any such error. Your suggestions will be appreciated.

EDITIONS

First Edition : 2020

Second Edition : 2021

Third Edition : 2022

Fourth Edition : 2023

Fifth Edition : 2024

Sixth Edition : 2025

Seventh Edition : 2026

CONTENTS

Engineering Mechanics

CHAPTER 1

Basics of Vectors 1 - 10

1.1	Vectors and Scalars.....	1
1.2	Equality of Vectors.....	1
1.3	Unit Vectors.....	2
1.4	Addition of Vectors.....	2
1.5	Multiplication of Vector by a number.....	3
1.6	Subtraction of Vectors.....	3
1.7	Resolution of Vectors.....	3
1.8	Dot Product or Scalar Product of Two Vectors.....	5
1.9	Dot product of two vectors in terms of the components along the coordinate axes.....	5
1.10	Cross product or Vector product of two vectors.....	6
1.11	Position Vector.....	7
1.12	Moment of a Force about a Point.....	8
1.13	Moment of Force about an Axis.....	8
1.14	Varignon's Theorem.....	8
	<i>Objective Brain Teasers</i>	9

CHAPTER 2

Motion in Space..... 11 - 23

2.1	Introduction.....	11
2.2	Distance and Displacement.....	11
2.3	Average speed and Instantaneous speed.....	11
2.4	Average velocity and Instantaneous velocity.....	12
2.5	Average acceleration and Instantaneous acceleration.....	12
2.6	Rectilinear motion.....	13
2.7	Curvilinear motion.....	15
	<i>Objective Brain Teasers</i>	21

CHAPTER 3

Composition, Resolution and Equilibrium of Forces..... 24 - 38

3.1	Introduction.....	24
3.2	Force Systems.....	25
3.3	Resultant Force.....	26
3.4	Composition of Forces.....	28
3.5	Resolution of Forces.....	29
3.6	Equilibrium of Forces.....	30
3.7	Principle of Transmissibility.....	33
3.8	Free Body Diagram.....	34
3.9	Newton's Laws of Motion.....	34
	<i>Objective Brain Teasers</i>	35
	<i>Conventional Brain Teasers</i>	37

CHAPTER 4

Friction 39 - 79

4.1	Introduction.....	39
4.2	Types of Friction.....	39
4.3	Laws of Friction.....	39
4.4	Coefficient of Friction.....	40
4.5	Angle of Friction.....	40
4.6	Angle of Repose.....	41
4.7	Force Analysis of a Sliding Body.....	41
4.8	Wedge.....	45
4.9	Rolling Friction.....	48
4.10	Motion of Vehicles on rough Road.....	48
4.11	Belt, Rope and Pulley.....	56

4.12 Two Mating Blocks	67
4.13 Friction in Wheels	68
<i>Objective Brain Teasers</i>	73
<i>Conventional Brain Teasers</i>	76

CHAPTER 5

Center of Gravity and Moment of Inertia 80 – 95

5.1 Introduction	80
5.2 Centroid	80
5.3 Centroid of Given Lamina.....	80
5.4 Moment of Inertia	85
<i>Objective Brain Teasers</i>	93
<i>Conventional Brain Teasers</i>	94

CHAPTER 6

Concept of Virtual Work, Energy and Power 96 - 114

6.1 Introduction	96
6.2 Principle of Virtual Work.....	96
6.3 Work and Energy.....	99
6.4 Rolling Motion under Gravity.....	102
6.5 Power	103
<i>Objective Brain Teasers</i>	108
<i>Conventional Brain Teasers</i>	110

CHAPTER 7

Impulse, Momentum and Rotational Motion 115 - 140

7.1 Momentum.....	115
7.2 Impulse.....	115
7.3 Impulse-Momentum Theorem	115
7.4 Law of Conservation of Linear Momentum	116
7.5 Elastic and Inelastic Impact.....	116
7.6 Rotational Motion	123
7.7 Rotational kinematics	123
7.8 Relation between linear motion and rotational motion of a particle on a body	124
7.9 Torque of a Force	125
7.10 Body in Equilibrium	127
7.11 Angular momentum.....	127
7.12 Mass moment of inertia	128
7.13 Rolling Without sliding or pure rolling	130
<i>Objective Brain Teasers</i>	133
<i>Conventional Brain Teasers</i>	136



Basics of Vectors

1.1 VECTORS AND SCALARS

Many physical quantities are completely described by a numerical value alone and are added according to the ordinary rules of algebra. As an example the mass of a system is described by saying that it is 5 kg. If two bodies one having a mass of 5 kg and the other having a mass of 2 kg are added together to make a composite system, the total mass of the system becomes $5 \text{ kg} + 2 \text{ kg} = 7 \text{ kg}$. Such quantities are called scalars.

So, a scalar is any positive or negative physical quantity that can be completely specified by its magnitude. Other examples of scalar quantities are length, mass and time.

While the complete description of certain physical quantities requires a numerical value as well as a direction in space. Velocity of a particle is an example of this kind. The magnitude of velocity is represented by a number such as 5 m/s and tells us how fast a particle is moving. But the description of velocity becomes complete only when the direction of velocity is also specified. We can represent this velocity by drawing a line parallel to the velocity and putting an arrow showing the direction of velocity.




Fig. Triangle law to obtain resultant velocity

Further, if a particle is given two velocities simultaneously, its resultant velocity is different from the two velocities and is obtained by using a special rule known as triangle law.

The physical quantities which have magnitude and direction and which can be added according to the laws of vector addition are called vector quantities. Other examples of vector quantities are force, linear momentum, electric field, magnetic field etc.

1.2 EQUALITY OF VECTORS

Two vectors (representing two values of the same physical quantity) are called equal if their magnitudes and directions are same. Thus, a parallel translation of a vector does not bring about any change in it.

1.3 UNIT VECTORS

A vector whose magnitude is unity (i.e. 1 unit) is called a unit vector. Unit vector is a director vector. Unit

vector in the direction of any given vector \vec{a} is given as $\frac{\vec{a}}{|\vec{a}|}$.

EXAMPLE : 1.1

The unit vector in the direction of $\vec{A} = 5\hat{i} + \hat{j} - 2\hat{k}$ is

Solution:

$$|\vec{A}| = \sqrt{5^2 + 1^2 + (-2)^2} = \sqrt{30}$$

$$\text{The required unit vector is } \frac{\vec{A}}{|\vec{A}|} = \frac{5}{\sqrt{30}}\hat{i} + \frac{1}{\sqrt{30}}\hat{j} - \frac{2}{\sqrt{30}}\hat{k}$$

1.4 ADDITION OF VECTORS

The triangle rule of vector addition is already described above. If \vec{a} and \vec{b} are two vectors to be added, a diagram is drawn in which the tail of \vec{b} coincides with the head of \vec{a} . The vector joining the tail of \vec{a} with the head of \vec{b} is the vector sum of \vec{a} and \vec{b} .

The same rule may be stated in a slightly different way by parallelogram law.




Fig. Vector addition as per triangle law

In parallelogram law we draw vectors \vec{a} and \vec{b} with both the tails coinciding as shown in figure. Taking these two as the adjacent sides we complete the parallelogram. The diagonal through the common tails gives the sum of the two vectors.

Suppose the magnitude of $\vec{a} = a$ and that of $\vec{b} = b$. If the angle between \vec{a} and \vec{b} is θ , then

$$\begin{aligned} AD^2 &= (AB + BE)^2 + (DE)^2 \\ &= (a + b\cos\theta)^2 + (b\sin\theta)^2 \\ &= a^2 + 2ab\cos\theta + b^2 \end{aligned}$$

Thus, the magnitude of $\vec{a} + \vec{b}$ is

$$= \sqrt{a^2 + b^2 + 2ab\cos\theta}$$

Its angle with \vec{a} is α where,

$$\tan\alpha = \frac{DE}{AE} = \frac{b\sin\theta}{a + b\cos\theta}$$




Fig. Vector addition as per parallelogram law

Special cases:

(a) When two vectors are acting in same direction, then,

$$\theta = 0^\circ$$

\therefore

$$|\vec{a} + \vec{b}| = \sqrt{a^2 + b^2 + 2ab} = a + b$$

and
$$\tan\alpha = \frac{b \times \sin 0^\circ}{a + b \cos 0^\circ} = 0$$

$\Rightarrow \alpha = 0^\circ$

Thus, the magnitude of sum of vectors \vec{a} and \vec{b} is equal to the sum of magnitudes of two vectors acting in same direction and their resultant acts in direction of \vec{a} and \vec{b} .

(b) When two vectors acts in opposite directions:

Then,

$$\theta = 180^\circ$$

\therefore

$$|\vec{a} + \vec{b}| = \sqrt{a^2 + b^2 - 2ab} = a - b$$

and

$$\tan\alpha = \frac{b \times \sin(180^\circ)}{a + b \times \cos(180^\circ)} = 0$$

\Rightarrow

$$\alpha = 0^\circ \text{ or } 180^\circ$$

Thus, the magnitude of sum of the vectors \vec{a} and $(-\vec{b})$ is equal to the difference of magnitudes of two vectors and their resultant acts in direction of bigger vector.

1.5 MULTIPLICATION OF VECTOR BY A NUMBER

Suppose \vec{a} is a vector of magnitude a and k is a number. We define the vector $\vec{b} = k\vec{a}$ as a vector of magnitude $|ka|$. If k is positive the direction of the vector $\vec{b} = k\vec{a}$ is same as that of \vec{a} . If k is negative, the direction of \vec{b} is opposite to \vec{a} . In particular, multiplication by (-1) just inverts the direction of the vector. The vectors \vec{a} and $-\vec{a}$ have equal magnitudes but opposite directions.

If \vec{a} is a vector of magnitude a and \hat{u} is a vector of unit magnitude in the direction of \vec{a} , we can write $\vec{a} = a\hat{u}$.

1.6 SUBTRACTION OF VECTORS

Let \vec{a} and \vec{b} be two vectors. We define $\vec{a} - \vec{b}$ as the sum of the vector \vec{a} and the vector $(-\vec{b})$. To subtract \vec{b} from \vec{a} , invert the direction of \vec{b} and add to \vec{a} .




Fig. Vector diagram representing $\vec{a} + (-\vec{b})$

1.7 RESOLUTION OF VECTORS

Consider a vector $\vec{a} = \vec{OA}$ in the X-Y plane drawn from the origin O. The length OB is called the projection of \vec{OA} on X-axis. Similarly OC is the projection of \vec{OA} on Y-axis. According to the rules of vector addition

$$\vec{a} = \vec{OA} = \vec{OB} + \vec{OC}$$

We have resolved the vector \vec{a} into two parts, one along OX and the other along OY. The magnitude of the part along OX is $OB = a \cos\alpha$ and the magnitude of the part along OY is $OC = a \cos\beta$. If \hat{i} and \hat{j} denote vectors of unit magnitude along OX and OY respectively, we get

$$\vec{OA} = a \cos\alpha \hat{i} \text{ and } \vec{OC} = a \cos\beta \hat{j}$$

So, that,

$$\vec{a} = a \cos\alpha \hat{i} + a \cos\beta \hat{j}$$

If the vector \vec{a} is not in the X-Y plane, it may have nonzero projections along X, Y, Z axes and we can resolve it into three parts i.e. along the X, Y and Z axes. If α, β, γ be the angles made by the vector \vec{a} with the three axes respectively, we get

$$\vec{a} = a \cos\alpha \hat{i} + a \cos\beta \hat{j} + a \cos\gamma \hat{k}$$

where \hat{i} , \hat{j} and \hat{k} are the unit vectors along X, Y and Z axes respectively. The component of vector \vec{a} along direction making angle θ with it is $a \cos\theta$ which is the projection of \vec{a} along the given direction.




Fig. Resolution of vector OA in x and y directions




Fig. Horizontal projection of \vec{a}

EXAMPLE : 1.2

The magnitudes of vectors \vec{OA} , \vec{OB} and \vec{OC} in the figure shown are equal. Find the direction of $\vec{OA} + \vec{OB} - \vec{OC}$.

Solution:

Let,

$$OA = OB = OC = F$$

$$x\text{-component of } \vec{OA} = F \cos 30^\circ = F \frac{\sqrt{3}}{2}$$


$$x\text{-component of } \vec{OB} = F \cos 60^\circ = \frac{F}{2}$$

$$x\text{-component of } \vec{OC} = F \cos 135^\circ = -\frac{F}{\sqrt{2}}$$

$$x\text{-component of } \vec{OA} + \vec{OB} - \vec{OC} = \left(\frac{F\sqrt{3}}{2}\right) + \left(\frac{F}{2}\right) - \left(-\frac{F}{\sqrt{2}}\right) = \frac{F}{2}(\sqrt{3} + 1 + \sqrt{2})$$

$$y\text{-component of } \vec{OA} = F \cos 60^\circ = \frac{F}{2}$$

$$y\text{-component of } \vec{OB} = F \cos 120^\circ = -\frac{F\sqrt{3}}{2}$$



$$y\text{-component of } \overline{OC} = F \cos 45^\circ = \frac{F}{\sqrt{2}}$$

$$y\text{-component of } \overline{OA} + \overline{OB} - \overline{OC} = \left(\frac{F}{2}\right) + \left(-\frac{F\sqrt{3}}{2}\right) - \left(\frac{F}{\sqrt{2}}\right) = \frac{F}{2}(1 - \sqrt{3} - \sqrt{2})$$

Angle of $\overline{OA} + \overline{OB} - \overline{OC}$ with the x -axis

$$= \tan^{-1} \frac{\frac{F}{2}(1 - \sqrt{3} - \sqrt{2})}{\frac{F}{2}(1 + \sqrt{3} + \sqrt{2})} = \tan^{-1} \frac{(1 - \sqrt{3} - \sqrt{2})}{(1 + \sqrt{3} + \sqrt{2})}$$

1.8 DOT PRODUCT OR SCALAR PRODUCT OF TWO VECTORS

The dot product (also called scalar product) of two vectors \vec{a} and \vec{b} is defined as

$$\vec{a} \cdot \vec{b} = ab \cos \theta$$

The dot product is commutative and distributive.

$$\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$$

$$\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$$




Fig. Vector \vec{a} and \vec{b} having angle θ between them

EXAMPLE : 1.3

The work done by a force \vec{F} during a displacement \vec{r} is given by $\vec{F} \cdot \vec{r}$. Suppose a force of 12 N acts on a particle in vertically upward direction and particle is displaced through 2 m in the vertically downward direction. Find the work done by the force during this displacement?

Solution:

The angle between the force \vec{F} and the displacement \vec{r} is 180° . Thus, the work done is

$$W = \vec{F} \cdot \vec{r} = Fr \cos \theta = 12 \times 2 \times \cos 180^\circ$$

$$W = -24 \text{ N-m} = -24 \text{ J}$$

1.9 DOT PRODUCT OF TWO VECTORS IN TERMS OF THE COMPONENTS ALONG THE COORDINATE AXES

Consider two vectors \vec{a} and \vec{b} represented in terms of the unit vectors $\hat{i}, \hat{j}, \hat{k}$ along the coordinate axes as

$$\vec{a} = a_x \hat{i} + a_y \hat{j} + a_z \hat{k}$$

and

$$\vec{b} = b_x \hat{i} + b_y \hat{j} + b_z \hat{k}$$

Then,

$$\vec{a} \cdot \vec{b} = (a_x \hat{i} + a_y \hat{j} + a_z \hat{k}) \cdot (b_x \hat{i} + b_y \hat{j} + b_z \hat{k})$$

$$= \left[\begin{array}{l} a_x b_x \hat{i} \cdot \hat{i} + a_x b_y \hat{i} \cdot \hat{j} + a_x b_z \hat{i} \cdot \hat{k} \\ + a_y b_x \hat{j} \cdot \hat{i} + a_y b_y \hat{j} \cdot \hat{j} + a_y b_z \hat{j} \cdot \hat{k} \\ + a_z b_x \hat{k} \cdot \hat{i} + a_z b_y \hat{k} \cdot \hat{j} + a_z b_z \hat{k} \cdot \hat{k} \end{array} \right] \quad \dots (1)$$



**OBJECTIVE
BRAIN TEASERS**

Q.1 A particle whose speed is 25 m/sec moves along the line from A(2, 1) to B(9, 25). The velocity vector of the particle in the form $a\hat{i} + b\hat{j}$ is

- (a) $(7\hat{i} + 7\hat{j})$ m/s (b) $(7\hat{i} + 24\hat{j})$ m/s
(c) $(24\hat{i} + 12\hat{j})$ m/s (d) $(24\hat{i} + 7\hat{j})$ m/s

Q.2 Two forces are acting on a body, $\vec{F}_1 = 2\hat{i} + 3\hat{j}$ and it does 8J of work, $\vec{F}_2 = 3\hat{i} + 5\hat{j}$ and it does -4J of work on body. The magnitude of displacement traversed by the body is
(a) 57.27 m (b) 30.53 m
(c) 54.40 m (d) 61.06 m

Q.3 The angle between two vectors $\vec{A} = 2\hat{i} + \hat{j} - \hat{k}$ and $\vec{B} = \hat{i} - \hat{k}$ is
(a) 30° (b) 60°
(c) 0° (d) 90°

Q.4 If two vectors are given as:
 $\vec{A} = \hat{i} + 2\hat{j} - \hat{k}$
 $\vec{B} = -\hat{i} - \hat{j} + \hat{k}$

then, the vector perpendicular to $(\vec{A} \times \vec{B})$ can be

- (a) $2\hat{i} + 4\hat{j} + 2\hat{k}$ (b) $\hat{i} + \hat{j} + \hat{k}$
(c) $25\hat{i} - 625\hat{j} - 25\hat{k}$ (d) $3\hat{i} - 2\hat{j} + 3\hat{k}$


Q.5 If $(\vec{a} + \vec{b})$ is perpendicular to \vec{b} and $\vec{a} + 2\vec{b}$ is perpendicular to \vec{a} . If $|\vec{a}| = a$ and $|\vec{b}| = b$, then
(a) $a = b$ (b) $a = 2b$
(c) $b = 2a$ (d) $a = b\sqrt{2}$

ANSWER KEY

1. (b) 2. (d) 3. (a) 4. (c) 5. (d)

HINTS & EXPLANATIONS

1. (b)
Velocity vector is given by the product of magnitude of velocity (speed) multiplied by the unit vector along velocity.



Unit vector along velocity,

$$\widehat{AB} = \frac{\overline{AB}}{|AB|} = \frac{(9-2)\hat{i} + (25-1)\hat{j}}{\sqrt{7^2 + 24^2}}$$

$$\begin{aligned} \vec{v} &= |\vec{v}| \widehat{AB} = 25 \left(\frac{7\hat{i} + 24\hat{j}}{25} \right) \\ &= (7\hat{i} + 24\hat{j}) \text{ m/s} \end{aligned}$$

2. (d)
Let the displacement of the body is given by

$$\vec{r} = x\hat{i} + y\hat{j}$$

work done by first force,

$$W_1 = \vec{F}_1 \cdot \vec{r} = 8$$

$$\therefore (2\hat{i} + 3\hat{j}) \cdot (x\hat{i} + y\hat{j}) = 8$$

$$2x + 3y = 8 \quad \dots (i)$$

Work done by second force,

$$W_2 = \vec{F}_2 \cdot \vec{r} = -4$$