

18 *Years*

Previous Years Solved Papers

Civil Services Main Examination

(2009-2026)

Mathematics Paper-I

Topicwise Presentation



**MADE EASY Publications Pvt. Ltd.**

Corporate Office: 44-A/4, Kalu Sarai (Near Hauz Khas Metro Station), New Delhi-110016

E-mail: infomep@madeeasy.in

Contact: 9021300500

Visit us at: www.madeeasypublications.org

Civil Services Main Examination Previous Solved Papers : Mathematics Paper-I

© Copyright, by MADE EASY Publications Pvt. Ltd.

All rights are reserved. No part of this publication may be reproduced, stored in or introduced into a retrieval system, or transmitted in any form or by any means (electronic, mechanical, photo-copying, recording or otherwise), without the prior written permission of the above mentioned publisher of this book.

First Edition: 2020

Second Edition: 2021

Third Edition: 2022

Fourth Edition: 2023

Fifth Edition: 2024

Sixth Edition: 2025

Seventh Edition: 2026

Eighth Edition: 2027

Preface

Civil Service is considered as the most prestigious job in India and it has become a preferred destination by all engineers. In order to reach this estimable position every aspirant has to take arduous journey of Civil Services Examination (CSE). Focused approach and strong determination are the pre-requisites for this journey. Besides this, a good book also comes in the list of essential commodity of this odyssey.



B. Singh (Ex. IES)

I feel extremely glad to launch the fourth edition of such a book which will not only make CSE plain sailing, but also with 100% clarity in concepts.

MADE EASY team has prepared this book with utmost care and thorough study of all previous years papers of CSE. The book aims to provide complete solution to all previous years questions with accuracy.

I would like to acknowledge efforts of entire MADE EASY team who worked day and night to solve previous years papers in a limited time frame and I hope this book will prove to be an essential tool to succeed in competitive exams and my desire to serve student fraternity by providing best study material and quality guidance will get accomplished.

With Best Wishes

B. Singh (Ex. IES)

CMD, MADE EASY Group

Previous Years Solved Papers of
Civil Services Main Examination

Mathematics : Paper-I

C O N T E N T S

Sl.	TOPIC	PAGE No.
Unit-1	Linear Algebra.....	1 - 86
	1. Vector Space Over R and C	1
	2. Linear Transformations	17
	3. Matrix	71
Unit-2	Calculus.....	87 - 156
	1. Function of a Real Variable	87
	2. Functions of Two and Three Variables	108
	3. Riemann's Definition of Definite Integrals	134
	4. Indefinite Integrals	140
	5. Infinite and Improper Integrals.....	140
	6. Double and Triple Integrals	142
	7. Area, Surface and Volume.....	149
Unit-3	Analytic Geometry	157 - 221
	1. Straight Lines.....	157
	2. Shortest Distance between Two Skew Lines.....	161
	3. Plane and its Properties	164
	4. Sphere and its Properties	173
	5. Cone and its Properties.....	187
	6. Paraboloid and its Properties	201
	7. Ellipsoid and its Properties	211
	8. Hyperboloid of One and Two Sheets and its Properties.....	214

Unit-4	Ordinary Differential Equations	222 - 317
1.	Formulation of Differential Equations	222
2.	Equation of 1st Order and 1st Degree	225
3.	Equation of 1st Order but NOT of 1st Degree	251
4.	Second and Higher Order Linear Equation with Constant Coefficients	258
5.	Wronskian	272
6.	Second Order Linear Equations with Variable Coefficient	273
7.	Determination of Complete Solution when one solution is known and Method of Variation of Parameter	283
8.	Laplace and Inverse Laplace Transformation and Properties	299
9.	Application to Initial Value Problem for 2nd Order Linear Equation with Constant Coefficient	315
Unit-5	Dynamics & Statics	318 - 399
1.	Rectilinear Motion	318
2.	Motion in a Plane	329
3.	Constrained Motion	351
4.	Work & Energy	357
5.	Equilibrium of a System of Particles	366
6.	Work and Potential Energy	389
7.	Principle of Virtual Work	396
Unit-6	Vector Analysis	400 - 471
1.	Scalar and Vector Fields	400
2.	Differentiation of a Vector Field of a Scalar Variable	401
3.	Gradient, Divergence and Curl in Cartesian and Cylindrical Coordinate and Directional Derivative	405
4.	Vector Identity and Vector Equations	415
5.	Application to Geometry	421
6.	Curves in Space	426
7.	Gauss-Divergence Theorem, Stokes Theorem and Green's Identity	436



1. Vector Space Over R and C

- 1.1 Prove that the set V of the vectors (x_1, x_2, x_3, x_4) in R^4 which satisfy the equations $x_1 + x_2 + 2x_3 + x_4 = 0$ and $2x_1 + 3x_2 - x_3 + x_4 = 0$ is a subspace of R^4 . What is the dimension of this subspace. Find one of its bases.

(2009 : 12 Marks)

Solution:

ILD for vertical reaction at A;

$$V = \{(x_1, x_2, x_3, x_4) \in R^4 \mid x_1 + x_2 + 2x_3 + x_4 = 0, 2x_1 + 3x_2 - x_3 + x_4 = 0\}$$

Then clearly $(0, 0, 0, 0) \in V$ and so V is non-empty.Again let $x = (x_1, x_2, x_3, x_4) \in V$ and $y = (y_1, y_2, y_3, y_4) \in V$. Also let $\alpha, \beta \in R$.

$$\alpha x + \beta y = (\alpha x_1 + \beta y_1, \alpha x_2 + \beta y_2, \alpha x_3 + \beta y_3, \alpha x_4 + \beta y_4) \in R^4$$

Also, $\alpha(x_1 + x_2 + 2x_3 + x_4) + \beta(y_1 + y_2 + 2y_3 + y_4) = 0$ $\Rightarrow \alpha x + \beta y$ satisfies $x_1 + x_2 + 2x_3 + x_4 = 0$ Similarly $\alpha x + \beta y$ satisfies 2nd equation as well. $\therefore \alpha x + \beta y \in V$ **Dimension of the Subspace :**Any element of V is a solution to equation

$$\begin{bmatrix} 1 & 1 & 2 & 1 \\ 2 & 3 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

So, its dimension is same as rowspace of coefficient matrix, i.e., its rank.

Reducing it to row reduced echelon form

$$\begin{bmatrix} 1 & 1 & 2 & 1 \\ 2 & 3 & -1 & 1 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_2 - 2R_1} \begin{bmatrix} 1 & 1 & 2 & 1 \\ 0 & 1 & -5 & -1 \end{bmatrix}$$

As it has two non-zero rows in row reduced form.

$$\dim(V) = \text{Rank of matrix} = 2$$

Writing the equation as matrix

$$\begin{bmatrix} 1 & 1 & 2 & 1 \\ 2 & 3 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \text{ or } Ax = 0.$$

So,

Clearly

Let

$$V = \{x = (x_1, x_2, x_3, x_4) \in R^4 \mid Ax = 0\}$$

$$O = (0, 0, 0, 0) \in V \text{ so } V \text{ is non-empty.}$$

$$x = (x_1, x_2, x_3, x_4); y = (y_1, y_2, y_3, y_4) \in V \text{ and } \alpha, \beta \in R$$

$$A(\alpha x + \beta y) = A(\alpha x) + A(\beta y)$$

$$= \alpha(Ax) + \beta(Ay) = \alpha \cdot 0 + \beta \cdot 0 = 0$$

$$\therefore \alpha x + \beta y \in V$$

$\therefore V$ is a vector subspace.

Clearly, V is the null space of the matrix

$$A = \begin{bmatrix} 1 & 1 & 2 & 1 \\ 2 & 3 & -1 & 1 \end{bmatrix}$$

Reducing it to row reduced echelon form

$$\begin{bmatrix} 1 & 1 & 2 & 1 \\ 2 & 3 & -1 & 1 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_2 - 2R_1} \begin{bmatrix} 1 & 1 & 2 & 1 \\ 0 & 1 & -5 & -1 \end{bmatrix}$$

$$\dim(\text{Row space}(A)) = \text{Number of non-zero rows} \\ = 2$$

By Rank-Nullity Theorem :

$$\underset{\text{(Nullity)}}{\dim(\text{Null space})} + \underset{\text{(Rank)}}{\dim(\text{Row space})} = n = 4$$

$$\dim(\text{null space}) = 2$$

$$\dim(V) = 2$$

$\therefore$

For Finding Basis :

$$\dim(\text{null space}) = 2$$

$\therefore$

$$\text{No. of free variables} = 4 - 2 = 2$$

So, we fix 2 variables.

Taking $x_3 = 1, x_4 = 0$ first.

$$\begin{cases} x_1 + x_2 = -2 \\ 2x_1 + 3x_2 = 1 \end{cases} \Rightarrow \begin{cases} x_1 = -7 \\ x_2 = 5 \end{cases} \quad x = (-7, 5, 1, 0)$$

Taking $x_3 = 0, x_4 = 1$

$$\begin{cases} x_1 + x_2 = -1 \\ 2x_1 + 3x_2 = -1 \end{cases} \Rightarrow \begin{cases} x_1 = -2 \\ x_2 = 1 \end{cases} \quad x = (-2, 1, 0, 1)$$

$\therefore (-7, 5, 1, 0)$ and $(-2, 1, 0, 1)$ are two elements of V . And since they are linearly independent (**because of choice of 3rd and 4th element**) they form a basis.

- 1.2 Prove that set V of all 3×3 real symmetric matrices form a linear subspace of the space of all 3×3 real matrices. What is the dimension of this subspace? Find at least one of the bases for V .**

(2009 : 20 Marks)

Solution:

Approach : Use definition of subspaces for first part. For the 2nd impose conditions due to symmetry on the matrix.

Let V be subset of all 3×3 symmetric matrix.

Then $I_3 \in V$ so V is not empty. Again, let $A, B \in V$.

$\Rightarrow$

$$A = A^T$$

$$B = B^T \text{ (definition of symmetric)}$$

and $\alpha, \beta \in R$.

Then

$$\begin{aligned} (\alpha A + \beta B)^T &= (\alpha A)^T + (\beta B)^T \\ &= \alpha A^T + \beta B^T = \alpha A + \beta B \end{aligned}$$

$\therefore \alpha A + \beta B \in V$.

So, V is a vector subspace of the space of all 3×3 real matrices over R .

Dimension : Let
$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \in V$$

$$\Rightarrow A^T = A \Rightarrow \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & i \end{bmatrix}$$

i.e.,
$$b = d, c = g, h = f$$

$\therefore$
$$A = \begin{bmatrix} a & b & c \\ b & e & f \\ c & f & i \end{bmatrix}$$

Thus, any general element has 6 variables (instead of 9 for a 3×3 real matrix). So, dimension of V is 6.

Basis : Putting each of the variables as 1 and rest 0 gives us a basis, i.e.,

$$B = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \right\}$$

1.3 In the n -space R^n , determine whether or not the set $\{e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n, e_n - e_1\}$ is linearly independent.

(2010 : 10 Marks)

Solution:

Given the set is $\{e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n, e_n - e_1\}$.

$$\text{Let } a_1(e_1 - e_2) + a_2(e_2 - e_3) + \dots + a_{n-1}(e_{n-1} - e_n) + a_n(e_n - e_1) = 0$$

$$\Rightarrow e_1(a_1 - a_n) + e_2(a_2 - a_1) + e_3(a_3 - a_2) + \dots + e_n(a_n - a_{n-1}) = 0$$

As $e_1, e_2, \dots, e_n$ from basis of R^n , so they are linearly independent.

$$\therefore a_1 - a_n = 0 \quad \dots(1)$$

$$a_2 - a_1 = 0 \quad \dots(2)$$

$$a_3 - a_2 = 0 \quad \dots(3)$$

$$a_n - a_{n-1} = 0 \quad \dots(n)$$

$\therefore$ from eqn. (1) to (n) it can be deduced that

$$a_1 = a_2 = a_3 = \dots a_{n-1} = a_n$$

and not need to be zero.

$\therefore$ The given set is linearly dependent.

1.4 Show that the subspaces of $\mathbb{R}^3$ spanned by two sets of vectors $\{(1, 1, -1), (1, 0, 1) \text{ and } (1, 2, -3), (5, 2, 1)\}$ are identical. Also find the dimension of this subspace.

(2011 : 10 Marks)

Solution :

Let W_1 be the subspace generated by the vectors $(1, 1, -1), (1, 0, 1)$.

Consider a matrix A , whose rows are the given vectors $(1, 1, -1), (1, 0, 1)$ and reduce it to row reduced Echelon form.

$$\therefore A = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 0 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & -1 \\ 0 & -1 & 2 \end{bmatrix} \text{ by } R_2 \leftrightarrow R_2 - R_1 \quad \dots(i)$$

Again W_2 be the subspace generated by the vectors $(1, 2, -3), (5, 2, 1)$.

Consider a matrix B , whose rows are the given vectors $(1, 2, -3)$ and $(5, 2, 1)$ and reduce it to row reduced Echelon form.

$$\text{i.e., } B = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 2 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 2 & -3 \\ 0 & -8 & 16 \end{bmatrix} \text{ by } R_2 \rightarrow R_2 - 5R_1$$

$$\sim \begin{bmatrix} 1 & 2 & -3 \\ 0 & -1 & 2 \end{bmatrix} R_2 \rightarrow \frac{1}{8}R_2$$

$$\sim \begin{bmatrix} 1 & 1 & -1 \\ 0 & -1 & 2 \end{bmatrix} R_1 \rightarrow R_1 + R_2 \quad \dots(ii)$$

From (i) and (ii), we find that the non-zero rows in the row-reduced Echelon forms of matrices A and B are the same.

$$\therefore \text{Row space of } A = \text{Row space of } B$$

$$\therefore W_1 = W_2$$

Again consider the vectors $(1, 1, -1)$ and $(1, 0, 1)$.

$$\text{Let } a_1(1, 1, -1) + a_2(1, 0, 1) = (0, 0, 0), a_1, a_2, a_3 \in R$$

$$\Rightarrow (a_1, a_1, -a_1) + (a_2, 0, a_2) = (0, 0, 0)$$

$$\Rightarrow (a_1 + a_2, a_1, -a_1 + a_2) = (0, 0, 0)$$

$$\Rightarrow a_1 + a_2 = 0, a_1 = 0, -a_1 + a_2 = 0$$

$$\Rightarrow a_1 = 0 = a_2$$

$\Rightarrow$ The vectors $(1, 1, -1)$ and $(1, 0, 1)$ are linearly independent.

$\therefore$ The vectors form basis of $W_1 = W_2$.

$\therefore$ Dimension of $W_1 (= W_2) = 2$.

1.5 Prove or disprove the following statement :

If $B = \{b_1, b_2, b_3, b_4, b_5\}$ is a basis for R^5 and V is a two-dimensional subspace of R^5 , then V has a basis made of just two members of B .

(2012 : 12 Marks)

Solution :

$$B = \{b_1, b_2, b_3, b_4, b_5\} \text{ is a basis for } R^5.$$

V is a two-dimensional subspace of R^5 .

Consider the set $\{b_1, b_2\}$

$$\text{Let } B' = \{b_1, b_2\}$$

Since $B = \{b_1, b_2, b_3, b_4, b_5\}$ is a basis of R^5 .

$\Rightarrow b_1, b_2, b_3, b_4, b_5$ are linearly independent and any subset of a L.I. set of vectors is L.I. b_1, b_2 are L.I.

Also, if a basis of vector space contains n elements, then any subset of the vector space having n elements is a basis iff the subset is L.I.

Thus, V will have a basis made of just 2 members of B iff B' is a subset of V .

- 1.6 Let V be the vector space of all 2×2 matrices over the field of real numbers. Let W be the set consisting of all matrices with zero determinant. Is W a subspace of V ? Justify your answer.
(2012 : 8 Marks)

Solution:

Let $W_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, then $|W_1| = \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0$

and $W_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, then $|W_2| = \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 0$

Thus, $W_1, W_2 \in W$

Now, $W_1 + W_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

and $|W_1 + W_2| = \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 \neq 0$

$\therefore W_1 + W_2 \notin W$

$\Rightarrow W$ is not a subspace of V as we know that a non-empty subset W of a vector space $V(F)$ is a subspace of V iff W is closed under addition and scalar multiplication.

- 1.7 Let V be an n -dimensional vector space and $T : V \rightarrow U$ be an invertible linear operator. If $\beta = \{X_1, X_2, \dots, X_n\}$ is a basis of V , show that $\beta' = \{TX_1, TX_2, \dots, TX_n\}$ is also a basis of U .
(2013 : 8 Marks)

Solution :

Approach : An invertible linear transformation into the same vector space is one-one onto. Anyways linearity of T allows use to prove β' is linearly independent and spans V .

An invertible linear transformation from a vector space to itself is one-one and onto.

$\therefore T$ is one-one and onto.

To prove $\beta' = \{TX_1, \dots, TX_n\}$ is a basis, we need to show that it spans V and is linearly independent.

Spans V :

Let $x \in V$. Since T is invertible and so one-one there exists $y \in V$ such that

$$T(y) = x$$

Now $y \in V$ and β is a basis.

So, $\exists C_1, C_2, \dots, C_n \in F$ such that

$$\begin{aligned} y &= C_1X_1 + \dots + C_nX_n \\ T(y) &= T(C_1X_1 + \dots + C_nX_n) \\ x &= C_1TX_1 + \dots + C_nTX_n \end{aligned}$$

$\Rightarrow$

$$\therefore x \in \text{Span}\{TX_1, \dots, TX_n\}$$

β' is linearly independent.

Let it be linearly dependent, then $\exists C_1, C_2, \dots, C_n \in F$ not all zero such that

$$C_1TX_1 + \dots + C_nTX_n = 0$$

$$\Rightarrow T(C_1X_1 + \dots + C_nX_n) = 0$$

But T is invertible $\Rightarrow T$ is one-one.

$$\therefore \text{Kernel } T = \{0\}$$

$$\Rightarrow C_1X_1 + \dots + C_nX_n = 0$$

with not all of C_i 's zero.

But this means $X_1, X_2, \dots, X_n$ are not linearly independent, a contradiction.

$\therefore \beta'$ is linearly independent set and thus β' is a basis.

- 1.8 Show that the vector $X_1 = (1, 1 + i, i)$, $X_2 = (i, -i, 1 - i)$ and $X_3 = (0, 1 - 2i, 2 - i)$ in C_3 are linearly independent over the field of real numbers but are linearly dependent over the field of complex numbers.

(2013 : 8 Marks)

Solution :

Approach : Linear independence depends on the constants of the scalar fields over which the vector space is defined. For real constants the linear combination can not be made 0. But it can be for complex constants.

Let X_1, X_2, X_3 be three vectors over R .

Let $C_1, C_2, C_3 \in R$ such that

$$\begin{aligned} & C_1 X_1 + C_2 X_2 + C_3 X_3 = 0 \\ \Rightarrow & C_1(1, 1 + i, i) + C_2(i, -i, 1 - i) + C_3(0, 1, -2i, 2 - i) = 0 \\ \Rightarrow & [C_1 + C_2 i, (C_1 + C_3) + (C_1 - C_2 - 2C_3)i, C_2 + 2C_3 + (C_1 - C_2 - C_3)i] = 0 \\ \Rightarrow & C_1 + C_2 i = 0 \\ \Rightarrow & (C_1 + C_3) + (C_1 - C_2 - 2C_3)i = 0 \quad (\text{using first two terms}) \\ \Rightarrow & C_1 = 0, C_2 = 0, C_3 = 0 \end{aligned}$$

So, X_1, X_2, X_3 are linearly independent over R .

Linear dependence on C .

Let $i, -1, 1 \in C$ be the constants.

$$\begin{aligned} & = i(1, 1 + i, i) + (-1)(i, -i, 1 - i) + 1(0, 1 - 2i, 2 - i) \\ & = (i, i - 1, -1) + (-i, i, -1 + i) + (0, 1 - 2i, 2 - i) = (0, 0, 0) \end{aligned}$$

$\therefore X_1, X_2, X_3$ are linearly dependent on C .

- 1.9 Find one vector in R^3 which generates the intersection of V and W , where V is the xy -plane and w is the space generated by the vectors $(1, 2, 3)$ and $(1, -1, 1)$.

(2014 : 10 Marks)

Solution:

Let $R^3 = \{(x, y, z) / x, y, z \in IR\}$ be the given vector space.

Given that V is the xy -plane.

$$\therefore V = \left\{ \frac{(x, y, z) \in IR^3}{z = 0 : x, y \in IR} \right\} \quad \dots(i)$$

and given that w is the space generated by the vectors $(1, 2, 3)$ and $(1, -1, 1)$.

For this, we find a homogeneous system whose solution set W is generated by

$$\begin{aligned} S &= \{(1, 2, 3), (1, -1, 1)\} \\ W &= \{\alpha(1, 2, 3) + \beta(1, -1, 1); \alpha, \beta \in IR\} \\ &= \{(\alpha + \beta, 2\alpha - \beta, 3\alpha + \beta), \alpha, \beta \in IR\} \\ V &= \{(x, y, 0); \lambda, y \in IR\} \end{aligned}$$

Now for $V \cap W$

$$\begin{aligned} \Rightarrow & x = \alpha + \beta, y = 2\alpha - \beta \text{ and } 3\alpha + \beta = 0 \\ \Rightarrow & \beta = -3\alpha \\ \Rightarrow & x = -2\alpha, y = 5\alpha \text{ and } z = 0 \\ \therefore & V \cap W = \{-2\alpha, 5\alpha, 0; \alpha \in IR\} \end{aligned}$$

Clearly $V \cap W$ is spanned by $(-2, 5, 0) \in IR^3$.

- 1.10 Let v and w be the following subspaces of R^4 .

$$v = \{(a, b, c, d) : b - 2c + d = 0\}$$

and

$$w = \{(a, b, c, d) : a = d, b = 2c\}$$

Find a basis and the dimension of (i) v (ii) w (iii) $v \cap w$.

(2014 : 15 Marks)

Solution:

We observe that $(a, b, c, d) \in G$

$$\Rightarrow b - 2c + d = 0$$

$$\begin{aligned} \Rightarrow (a, b, c, d) &= (a, 5, c, 2, cb) \\ &= (a, 0, 0, 0) + (0, b, 0, -b) + (0, 0, c, 2c) \\ &= a(1, 0, 0, 0) + b(0, 1, 0, -1) + c(0, 0, 1, 2) \end{aligned}$$

This shows that every vector in v is a linear combination of the three linearly independent vectors $(1, 0, 0, 0)$, $(0, 1, 0, -1)$, $(0, 0, 1, 2)$

Thus, a basis of v is

$$A = \{(1, 0, 0, 0), (0, 1, 0, -1), (0, 0, 1, 2)\}$$

Hence, **$\dim v = 3$**

Now, $(a, b, c, d) \in w \Rightarrow$

$$\begin{aligned} \Rightarrow a &= d, b = 2c \\ \Rightarrow (a, b, c, d) &= (a, 2c, c, a) = (a, 0, 0, a) + (0, 2c, c, 0) \\ &= a(1, 0, 0, 1) + c(0, 2, 1, 0) \end{aligned}$$

which shows that w is generated by the linearly independent set $\{(1, 0, 0, 1), (0, 2, 1, 0)\}$

$$\therefore \text{A basis for } w \text{ is } = \{(1, 0, 0, 1), (0, 2, 1, 0)\}$$

and

$$\dim w = 2$$

$$(a, b, c, d) \in v \cap w \Rightarrow (a, b, c, d) \in v \text{ and } (a, b, c, d)$$

$$\Rightarrow b - 2c + d = 0, a = d, b = 2c$$

$$\Rightarrow (a, b, c, d) = (0, 2c, c, 0) = c(0, 2, 1, 0)$$

Hence a basis of $v \cap w$ is $\{0, 2, 1, 0\}$

and

$$\dim(v \cap w) = 1$$

1.11 The vectors $v_1 = (1, 1, 2, 4)$, $v_2 = (2, -1, -5, 2)$, $v_3 = (1, -1, -4, 0)$ and $v_4 = (2, 1, 1, 6)$ are linearly independent. Is it true? Justify your answer.

(2015 : 10 Marks)

Solution:

We form a matrix with these given vectors as rows and reduce it to row-echelon form to investigate its rank.

$$A = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 2 & -1 & -5 & 2 \\ 1 & -1 & -4 & 0 \\ 2 & 1 & 1 & 6 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & -2 & -6 & -4 \\ 0 & -1 & -3 & -2 \end{bmatrix} \begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - R_1 \\ R_4 \rightarrow R_4 - 2R_1 \end{array}$$

$$\sim \begin{bmatrix} 1 & 1 & 2 & 4 \\ 0 & -3 & -9 & -6 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} R_3 \rightarrow R_3 - \frac{2}{3}R_2 \\ R_4 \rightarrow R_4 - \frac{R_2}{3} \end{array}$$

Since $\text{rank}(A) = 2 < \text{Number of rows (vectors)}$

Hence, the vectors are not linearly independent.

1.12 Find the dimension of the subspace of R^4 , spanned by the set $\{(1,0,0,0), (0,1,0,0), (1,2,0,1), (0,0,0,1)\}$. Hence, find its basis.

(2015 : 12 Marks)

Solution :

We find the echelon form of the matrix formed by given vectors taking as rows.

$$\begin{aligned} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - R_1} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 - 2R_2} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ & \sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_4 \rightarrow R_4 - R_3} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

Hence,

$$\text{Rank of Matrix} = 3$$

$\therefore$

$$\text{Dimension of subspace} = 3$$

For the basis, we take the vectors from the original matrix which correspond the non-zero rows in the echelon form.

$\therefore$

$$\text{Basis} = \{(1,0,0,0), (0,1,0,0), (1,2,0,1)\}$$

1.13 If

$$W_1 = \{(x, y, z) : x + y - z = 0\}$$

$$W_2 = \{(x, y, z) : 3x + y - 2z = 0\}$$

$$W_3 = \{(x, y, z) : x - 7y + 3z = 0\}$$

then find $\dim(W_1 \cap W_2 \cap W_3)$ and $\dim(W_1 + W_2)$.

(2016 : 3 Marks)

Solution :

$$\text{Let } (x, y, z) \in W_1 \cap W_2 \cap W_3$$

$\therefore$

$$x + y - z = 0 \quad \dots(i)$$

$$3x + y - 2z = 0 \quad \dots(ii)$$

$$x - 7y + 3z = 0 \quad \dots(iii)$$

$\Rightarrow$

$$x + y - z = 0$$

$\Rightarrow$

$$x + y - z = 0$$

$$-2y + z = 0$$

$\Rightarrow$

$$-2y + z = 0$$

$\therefore$

$$y = \frac{z}{2}$$

$$x = \frac{z}{2}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} z/2 \\ z/2 \\ z \end{bmatrix} = z \begin{bmatrix} 1/2 \\ 1/2 \\ 1 \end{bmatrix}$$

$$\begin{aligned}
 &\therefore \dim(W_1 \cap W_2 \cap W_3) = 1 \\
 \text{Again } &\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2) \quad \dots(*) \\
 \dim(W_1 \cap W_2) \Rightarrow &x + y - z = 0 \\
 &3x + y - 2z = 0 \\
 \Rightarrow &2x - z = 0 \\
 &\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} z/2 \\ z/2 \\ z \end{bmatrix} = \begin{bmatrix} 1/2 \\ 1/2 \\ 1 \end{bmatrix} z \\
 &\therefore \dim(W_1 \cap W_2) = 1 \\
 (*) \text{ gives } &\dim(W_1 + W_2) = 2 + 2 - 1 = 3
 \end{aligned}$$

1.14 Suppose U and W are distinct four dimensional subspaces of a vector space V , where $\dim V = 6$. Find the possible dimensions of subspace $U \cap W$.

(2017 : 10 Marks)

Solution:

$$\begin{aligned}
 &\dim(U) = \dim(W) = 4 \\
 &U \text{ and } W \text{ are subspaces of } V. \\
 \text{We know, } &\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W) \\
 \therefore &\dim(U \cap W) = \dim(U) + \dim(W) - \dim(U + W) \\
 &= 8 - \dim(U + W) \quad \dots(i) \\
 &\text{Now, } U + W \text{ is a subspace of } V \text{ while } U \text{ and } W \text{ are subspaces of } U + W. \\
 \therefore &\text{Possible } \dim(U + W) = 4, 5, 6 \\
 \text{Hence, by (3)} &\text{Possible } \dim(U \cap W) = 4, 3, 2 \\
 &\text{But } \dim(U \cap W) \text{ cannot be 4 otherwise } U \cap W = U = W, \text{ but } U \text{ and } W \text{ are distinct.} \\
 \therefore &\dim(U \cap W) = 2 \text{ or } 3
 \end{aligned}$$

1.15 Express basic vectors $e_1 = (1, 0)$ and $e_2 = (0, 1)$ as combinations of $\alpha_1 = (2, -1)$ and $\alpha_2 = (1, 3)$.

(2018 : 10 Marks)

Solution:

$$\begin{aligned}
 \text{Let any vector, say } &(x, y) = a\alpha_1 + b\alpha_2 \\
 \therefore &(x, y) = (2a + b, -a + 3b) \\
 \text{Comparing LHS and RHS, we get}
 \end{aligned}$$

$$a = \frac{3x - y}{7}, \quad b = \frac{x + 2y}{7}$$

For $e_1 = (1, 0) : x = 1, y = 0$

$$\therefore a = \frac{3}{7}, \quad b = \frac{1}{7}$$

$$\text{So, } e_1 = \frac{3}{7}\alpha_1 + \frac{1}{7}\alpha_2$$

For $e_2 = (0, 1) : x = 0, y = 1$

$$\therefore a = -\frac{1}{7}, \quad b = \frac{2}{7}$$

So,
$$e_2 = -\frac{1}{7}\alpha_1 + \frac{2}{7}\alpha_2$$

$$\therefore e_1 = \frac{3\alpha_1 + \alpha_2}{7}, e_2 = \frac{-\alpha_1 + 2\alpha_2}{7}$$

1.16 Consider the set V of all $n \times n$ real magic squares. Show that V is a vector space over R . Give examples of two distinct 2×2 magic squares.

(2020 : 10 Marks)

Solution:

Let V be the set of all $n \times n$ real magic squares. Let $\alpha, \beta \in V$, and sum of all rows and columns of α and β be k_1 and k_2 . Again sum of rows and column of matrix $\alpha + \beta$ will be $k_1 + k_2 \Rightarrow \alpha + \beta \in V$ (Internal composition). Now, let $a \in R$

Then $a\alpha \in V$ as the sum of rows and columns of $a\alpha$ will be ak_1 . Thus external composition is satisfied.

$\Rightarrow (V, +)$ forms an abelian group.

(i) $\alpha + \beta \in V$ for $\alpha, \beta, \gamma \in V$

(ii) $(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$

(iii) Zero matrix is identity

(iv) $-\alpha$ is inverse of α matrix

(v) Commutative property $\alpha + \beta = \beta + \alpha$. Again for $a, b \in R, \alpha, \beta \in V$, we have $a(\alpha + \beta) = a\alpha + b\beta$ (operation in matrix)

(vi) $(a + b)\alpha = a\alpha + b\beta$

(vii) $(ab)\alpha = a(b\alpha)$

(viii) $1 \cdot \alpha = \alpha$ (multiplication with identity matrix). Hence, V is a vector space. For examples: $\alpha = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix}$.

1.17 Prove that any set of n linearly independent vectors in a vector space V of dimension n constitutes a basis for V .

(2022 : 10 marks)

Solution:

Given $\dim V = n$

Let $S = \{a_1, a_2, a_3 \dots a_n\}$ be linearly independent (L.I.) set of n vectors in V .

If S is not a basis of V then it can be extended to form a basis of V .

In this case, the basis will contain more than n vectors.

But every basis of V must contain exactly n vectors.

Therefore our presumption is wrong and S must be basis of V .

1.18 Let the set $P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} : \begin{matrix} x - y - z = 0 \\ 2x - y + z = 0 \end{matrix} \right\}$ be the collection of vectors of a vector space $R^3(R)$. Then

(i) prove that P is a subspace of R^3 .

(ii) find a basis and dimension of P

(2022 : 10 + 10 marks)

Solution:

$$\text{Given } P = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} : x - y - z = 0 \text{ and } 2x - y + z = 0 \right\}$$

(i) We have $x - y - z = 0$

then

$$x = y + z$$

...(i)

and $2x - y + z = 0$, thus the value of $x = y + z$ then we have

$$2(y + z) - y + z = 0$$

$$\Rightarrow 2y + 2z - y + z = 0$$

$$\Rightarrow y + 3z = 0 \Rightarrow y = -3z$$

then

$$x = -3z + z = -2z$$

Thus, we have $(x, y, z) = (-2z, -3z, z) = z(-2, -3, 1)$

So, we have $P = \{z(-2, -3, 1) \text{ for } z \in R\}$

To prove, P is the vector space of $R^3(R)$

Let

$$\alpha_1 = z_1(-2, -3, 1) \text{ and } \alpha_2 = z_2(-2, -3, 1)$$

belongs to P then $a\alpha_1 + b\alpha_2 \in P$

where $\alpha_1, \alpha_2 \in P$ and $a, b \in R$

we have, $a\{z_1(-2, -3, 1)\} + b\{z_2(-2, -3, 1)\}$

$$= \{-2(az_1 + bz_2), -3(az_1 + bz_2), (az_1 + bz_2)\} \in P \text{ as } az_1 + bz_2 \in R$$

Thus P is the vector space of $R^3(R)$. Hence proved.

(ii) As we know $P = \{z(-2, -3, 1) \text{ for all } z \in R\}$

Thus $\text{Dim}(P) = 1$

and

$$\text{Basis set of } P = \begin{bmatrix} -2 \\ -3 \\ 1 \end{bmatrix}$$

1.19 Find a linear map $T: R^2 \rightarrow R^2$ which rotates each vector of R^2 by an angle θ . Also, prove that for

$\theta = \frac{\pi}{2}$, T has no eigenvalue in R .

(2022 : 15 marks)

Solution:

Let $A(x, y)$ is the initial vector of R^2 and after rotation by an angle θ the final vector becomes $B(x_1, y_1)$.


Here, $x = r \cos \phi$, $y = r \sin \phi$... (i)

$\therefore$

$$(x, y) = (r \cos \phi, r \sin \phi)$$

The final vector (x_1, y_1) after a rotation by angle θ becomes

$$\begin{aligned} x_1 &= r \cos(\theta + \phi) \\ &= r[\cos \theta \cos \phi - \sin \theta \sin \phi] \\ &= r \cos \theta \cos \phi - r \sin \theta \sin \phi \end{aligned}$$



Put the value of x and y from eq.(i), we have

$$x_1 = x \cos \theta - y \sin \theta$$

Now, calculate y_1 , we have

$$\begin{aligned} y_1 &= r \sin(\theta + \phi) \\ &= r[\sin \theta \cos \phi + \cos \theta \sin \phi] = r \sin \theta \cos \phi + r \cos \theta \sin \phi \end{aligned}$$

Put the value of x and y from eq.(ii), we have $y_1 = x \sin \theta + y \cos \theta$

Thus, the final vector $(x_1, y_1) \in \mathbb{R}^2$ becomes

$$(x_1, y_1) = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$$

Now for $\theta = \frac{\pi}{2}$, we have to prove T has no eigen value.

Let's find R_θ by presenting (x_1, y_1) in matrix form

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$\downarrow$
 R_θ

Now Put $\theta = \frac{\pi}{2}$, we have

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

We have $R_{\pi/2} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

The characteristic equation will be

$$\begin{aligned} |R_{\pi/2} - \lambda I| &= \begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix} - \begin{vmatrix} \lambda & 0 \\ 0 & \lambda \end{vmatrix} \\ &= \begin{vmatrix} -\lambda & -1 \\ 1 & -\lambda \end{vmatrix} = 0 \end{aligned}$$

$$\Rightarrow \lambda^2 - (-1) = 0$$

$$\Rightarrow \lambda^2 + 1 = 0 \Rightarrow \lambda^2 = -1$$

$$\Rightarrow \lambda = \pm i \quad \text{(Complex number)}$$

Thus $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ for $R(\pi/2)$, we can't have any eigen value in Real field ($\mathbb{R}^2$). Hence proved.

1.20 Let $V_1 = (2, -1, 3, 2)$, $V_2 = (-1, 1, 1, -3)$ and $V_3 = (1, 1, 9, -5)$ be three vectors of the space $\mathbb{R}^4$. Does $(3, -1, 0, -1) \in \text{span}\{V_1, V_2, V_3\}$? Justify your answer.

(2023 : 10 marks)

Solution:

$$\begin{aligned} \text{Let,} \quad (3, -1, 0, -1) &= x(2, -1, 3, 2) + y(-1, 1, 1, -3) + z(1, 1, 9, -5) \\ &= (2x - y + z) - x + y + z, 3x + y + 9z, 2x - 3y - 5z \end{aligned}$$

$$\text{Now,} \quad 2x - y + z = 3$$